

# G Y P S Y M O T H



### Background

The gypsy moth is considered to be the most important defoliating insect of hardwood trees in the Eastern United States. A French naturalist who was trying to develop a hardy strain of silkworm introduced it into the Boston area from Europe in 1869. Several of the moths escaped in 1870 and within ten years, they had established a viable population in the local area. Lacking their native European natural control agents, they spread rapidly and by 1889, this insect had defoliated over 230,000 acres of forest and shade trees. Despite the expenditure of millions of dollars in control efforts, this pest has continued to spread and now has caused millions of acres of defoliation, primarily in the northeastern states.

### How did it get to Idaho?

The gypsy moth is a notorious hitchhiker. The female moth lays her eggs on any solid surface including outdoor furniture, recreational vehicles, firewood, toys, etc. When these objects are later moved, gypsy moths can be transported long distances. This has been the main source of gypsy moth introductions to Idaho.

### Why is the gypsy moth a serious pest?

Gypsy moth caterpillars will feed on more than 300 kinds of plants. Preferred hosts in Idaho include paper birch, hawthorn, plum, oaks, apple, alder, aspen, filbert, and willow. Recent studies have also found that it can survive and reproduce on Douglas-fir, western larch, lodgepole pine, and other western conifers.

This insect has a tremendous capacity to increase in numbers, thus populations can rapidly build to large infestations causing widespread defoliation, which weakens, and sometimes kills trees. Weakened trees often become the targets of other killing insects and diseases.

Defoliation also reduces the aesthetic, recreational, and economic value of forests, parks, and wooded homesites. It can also impact stream side vegetation contributing to degradation of water quality.

The greatest economic threat to Idaho comes from the potential for newly infested states to be subject to restrictive quarantines to prevent further spread of the insect. Thus, our nursery, Christmas tree, and lumber industries could be heavily impacted.

When gypsy moth caterpillars are very numerous, they can be a nuisance to homeowners, crawling over buildings, vehicles, roads, and lawn furniture. Some people suffer allergic reactions when they contact hairs from the caterpillar.

### Life Cycle

The gypsy moth goes through four life stages: **egg**, **larva** (caterpillar), **pupa**, and **adult moth**. It has one generation per year and overwinters in the egg stage. Each female lays 50 to 1,000 **eggs** in one mass, which is covered with velvety golden or buff-colored hairs from the female's abdomen. The egg mass is about 3/4 inch wide and 1 to 1 1/5 inches long and may be attached to trees (Figure 1), logs, rocks, buildings, toys, or on outdoor household articles or vehicles.

Tom Schultz  
Director  
Idaho Department of Lands  
300 N. 6th Street, Suite 103  
Boise, ID 83720  
Phone: (208) 334-0200

**Insect and Disease**  
**No. 5**  
**September 2014**

Craig Foss  
Chief, Bureau of Forestry  
Assistance  
3284 W. Industrial Loop  
Coeur d'Alene, ID 83815  
Phone: (208) 769-1525

# GYPSY MOTH

**Caterpillars** hatch from eggs in mid-April to mid-June. When about half-grown, the caterpillar acquires five pairs of blue bumps and six pairs of red bumps on its back. These blue and red bumps distinguish the gypsy moth caterpillars from other kinds of hairy caterpillars. The full-grown caterpillar (Figure 2) is about 1 1/5 to 2 inches long.

In July, the caterpillar transforms into a non-feeding stage called the **pupa** (Figure 3). The adult moth develops inside this hard, dark-brown shell. The pupa is attached to solid objects by several strands of silk, often with the last skin shed by the caterpillar attached to its pointed end.

**Adult moths** begin to emerge in late July. The female moth (2-inch wing span) is white with brown zigzag markings on its wings (Figure 4). The male moth (Figure 5) is smaller (1 1/5-inch wing span) with mottled brown wings. It can be recognized by two feather-like antennae on its head. Adult moths do not feed. They live for about one week, during which time the sexes mate. Females lay eggs during August and early September starting the cycle over again.

## How can you help stop the gypsy moth?

1. Report suspected gypsy moth life stages to the Idaho Department of Lands.
2. Encourage anyone you know who has moved here recently from the northeastern U.S. to contact the Idaho Department of Lands for a **free inspection** of outdoor household articles and recreational vehicles.
3. Cooperate with the Idaho Department of Lands gypsy moth survey staff when they request permission to place traps each summer.
4. Help restrict movement of the gypsy moth by not moving wood products, firewood, plant material, outdoor household articles, or recreational vehicles out of gypsy moth infested areas without certification.

## Principal control techniques

The principal control techniques and/or tools currently available for use against the gypsy moth include chemical and biological pesticides, insect growth regulators (IGRs) and pheromones. Pheromones are synthetic reproductions of chemicals produced by moths to convey messages. Formulations of pheromone can be used as baits in traps for detection surveys or for control techniques such as mating disruption or mass trapping. For mating disruption the pheromone is formulated to be spread aerially to saturate the area "confusing" males. Similarly mass trapping saturates the area with traps, catching males before they are able to mate. Both of these techniques disrupt the mating cycle. Any of these techniques may be used independently or in conjunction with each other .

**Tree species** not favored by the gypsy moth include ash, true firs, cottonwood, catalpa, cedar, dogwood, sycamore, rhododendron, and tulip poplar.



Figure 1. *Gypsy moth egg mass.*



# GYPSY MOTH



Figure 2. *Full grown gypsy moth caterpillar.*



Figure 3. *Gypsy moth pupa.*




Figure 4. *Gypsy moth female.*



Figure 5. *Gypsy moth male.*



**FOR MORE INFORMATION CONTACT  
ANY IDAHO DEPARTMENT OF LANDS  
PRIVATE FORESTRY SPECIALIST**



Area	Office Location	Phone
Clearwater	Orofino	(208) 476-4587
Eastern Idaho	Idaho Falls	(208) 525-7167
Maggie Creek	Kamiah	(208) 935-2141
Mica	Coeur d'Alene	(208) 769-1577
Payette Lakes	McCall	(208) 634-7125
Pend Oreille Lake	Sandpoint	(208) 263-5104
Ponderosa	Deary	(208) 877-1121
Priest Lake	Coolin	(208) 443-2516
Southwest	Boise	(208) 334-3488
St Joe	St Maries	(208) 245-4551

<http://www.idl.idaho.gov/forestry/forest-health/index.html>